

## Research Article

# Efficiency of *Spirulina platensis* cultured with different nitrogen regimes on larval development, growth and survival rate of white shrimp, *Litopenaeus vannamei* n.R.<sup>1</sup>

Gorgij Jaski M.<sup>1\*</sup>; Rohani-Ghadikolaei K.<sup>2</sup>; Yahyavei M.<sup>1</sup>; Salarzadeh A.R.<sup>1</sup>

Received: August 2019

Accepted: May 2020

### Abstract

The blue-green algae *Spirulina platensis* was cultured with different nitrogen regimes (NH<sub>4</sub>Cl, NH<sub>4</sub>NO<sub>3</sub> and KNO<sub>3</sub>) with concentrations of 0.010, 0.025 and 0.050 M and then fed to *Litopenaeus vannamei* at 19 days post hatch (PL<sub>1</sub>) to evaluate the nutritional quality. At the end of the experiment, growth and survival rate were determined, and the larval development was calculated. The maximum protein content was obtained in culture media containing NH<sub>4</sub>NO<sub>3</sub> followed by NH<sub>4</sub>Cl and KNO<sub>3</sub> ( $p>0.05$ ). Moreover, in all *S. platensis* cultures, increases in nitrogen concentrations led to increases in protein content. The present study showed that, with the exception of shrimp larvae at 5 days post hatch, *L. vannamei* shrimp larvae fed on *S. platensis* cultured in NH<sub>4</sub>NO<sub>3</sub>, as a nitrogen regime, generally developed significantly ( $p<0.05$ ) faster to next stages compared with the other treatments. However, the survival rate (%) and total length (mm) of shrimp larvae fed on *S. platensis* did not show any significant difference ( $p>0.05$ ) when the nitrogen was varied in terms of regimes. Overall, the study points to the potential effectiveness of using NH<sub>4</sub>NO<sub>3</sub> as a nitrogen regime in *S. platensis* culture media in producing live food for the hatchery production of *L. vannamei* shrimp larvae.

**Keywords:** Nitrogen, *Spirulina platensis*, Survival rate, Larval development, *Litopenaeus vannamei*

---

1-Fisheries Department, Faculty of Natural Resources, Bandar Abbas Islamic Azad University, Bandar Abbas, Iran.

2-Persian Gulf and Oman Sea Ecology Research Center, Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research, Education and Extension Organization (AREEO), Bandar Abbas, Iran.REEO), Bandar Abbas, Iran.

\*Corresponding author's Email: sontderafshrimp@yahoo.com

## Introduction

The cost and composition of culture medium to biomass production of the microalgae are challenging factors for commercially viable production (Ilavarasi *et al.*, 2011). The first synthetic medium formulated for cultivation of *Spirulina* was Zarrouk's medium (Zarrouk, 1966) which is still used as the standard medium. Subsequently, different media have been tried for cultivation of *Spirulina* such as Rao's media (Singh, 2006), CFTIR media (Venkataraman *et al.*, 1995) and OFERR media (Singh, 2006). The production of *Spirulina* biomass requires water soluble forms of phosphorus and nitrogen (Mostert and Grobbelaar, 1987), and also sources of nutrition affect the growth rate and biochemical composition of *Spirulina*. The conventional nitrogen source for *S. platensis* is nitrate salts (sodium and potassium nitrates), and several studies have demonstrated the feasibility of replacing these conventional nitrogen sources with low-cost alternatives such as urea (Matsudo *et al.*, 2009; Avila-Leon *et al.*, 2012), ammonium sulfate (Ferreira *et al.*, 2010) and ammonium chloride (Bezerra *et al.*, 2008).

In the early stages of larval shrimp production, the supply of live food such as microalgae with high nutritional quality is one of the crucial activities in hatcheries (Spolaore *et al.*, 2006). Therefore, shrimp hatcheries are highly dependent on microalgae which are costly and labor intensive. Finding the efficient feeds for white shrimp larvae

is necessary in order to increase post-larvae quality and culture profitability. The dependence of commercial hatcheries on *Artemia* nauplii production will be a concern due to the possibility of rising cost of *Artemia* cyst, as well as the possible risk of disease transfer. Moreover, up to 80% of hatcheries have decreased using *Artemia* cyst in the last few years as a result of these concerns ((Lavens, and Sorgeloos, 2000)). In Iran, commercial hatcheries of *Litopenaeus vannamei* post larvae production need live food such as microalgae and *Artemia* nauplii as food at larval stages in their life cycle.

The major nutritional interest in *Spirulina* is due to high protein content, ease of digestion and significant contents of vitamins, minerals, amino acids and pigments (Piñero Estrada *et al.*, 2001; Marrez *et al.*, 2014; Kermani *et al.*, 2020). The cyanobacterium *Arthrospira (Spirulina) platensis* has even been used by humans because of its nutritional and possible medicinal effects (Colla *et al.*, 2007). Due to the fact that, *Spirulina* has many biological active substance, it has been used in aquaculture (Gouveia *et al.*, 2008) as well as poultry (Carrillo *et al.*, 2008) and ruminant (Kuplys *et al.*, 2009) nutrition. To date, a number of studies have been conducted using dried *Spirulina* as a feed supplement (Jin *et al.*, 2020; Holman and Malau-Aduli, 2012), while no study was conducted to use it in *L. vannamei* post larval production in Iran. Therefore, the main

objective of this study was to evaluate the performance of *L. vannamei* larvae (body length, survival and development index) fed on *S. platensis* cultured with different nitrogen regimes.

## Materials and methods

### *Microalgae culture*

*S. platensis* was grown in modified Zarrouk's medium (Rajasekaran *et al.*, 2016) in a 3000-ml Erlenmeyer flask, in which the  $\text{NaNO}_3$  was replaced by  $\text{NH}_4\text{Cl}$ ,  $\text{NH}_4\text{NO}_3$  and  $\text{KNO}_3$  with concentrations of 0.010, 0.025 and 0.050 M. Culture mediums were incubated at 30°C, salinity of 25 ppt and initial pH of 9.5 under 12:12 hour light-dark photoperiod with normal white light (Guillard and Ryther, 1962).

### *Protein analysis*

The *S. platensis* samples were harvested during the exponential growth phase (day 8) to determine protein content. Cells were concentrated using a centrifuge (Universal Tlettich, Germany) at 5000 g for 10 min and then freeze-dried (ZirBus, VaCo 5) before being stored at -25°C until biochemical analysis. Total protein was determined by the method of Lowry *et al.* (1951).

### *Larvae culture and feeding trial*

Newly hatched *L. vannamei* (nauplii 5;  $N_v$ ) larvae were supplied by a private shrimp hatchery (Sontderaf, Bandar Jask, Iran) where this study was conducted. For acclimation to the experimental conditions, the shrimp

nauplii were briefly kept in a 20-L tank filled with filtered and UV-treated seawater at 30°C and 35 mg  $\text{L}^{-1}$  salinity. At the start of the feeding trial, the shrimp nauplii were stocked in 5-L Erlenmeyer flasks filled with 3 L of filtered seawater at a density of 100 nauplii  $\text{L}^{-1}$  and reared for 19 days. The water was exchanged at a rate of 60–70% every other day. Three separate feeding regimes (consist of three replicates each) were conducted concurrently to evaluate their nutritional quality on early larval development, growth and survival of *L. vannamei* from nauplii 5 ( $N_v$ ) to postlarvae 9 ( $PL_{IX}$ ). The feeding regimes were as follows:

Larvae fed *S. platensis* that had been cultured with:

- (1)  $\text{NH}_4\text{Cl}$  (Diet A);
- (2)  $\text{NH}_4\text{NO}_3$  (Diet B); and
- (3)  $\text{KNO}_3$  (Diet C).

### *Larval development, total length and survival rate*

The early larval development, survival rate and total length of nauplii were evaluated at 5, 9, 14 and 19 days post hatch. The larval developmental stages were determined according to criteria defined by Ronquillo *et al.* (2006) for the larvae of *Penaeus semisulcatus*. The total length of individual shrimp larvae was measured to the nearest 0.01 mm by random sampling of 10 larvae from each dietary regime and under a binocular microscope from the tip of the rostrum to the end of the telson using an electronic digital caliper. The

mean survival rate (%) was calculated on given days by counting all survived larvae using glass pipettes, dividing by the total number that was initially stocked (300), and multiplying by 100.


#### Statistical analysis

Data were analyzed with SPSS 16.0 software using parametric tests and statistical analysis. A one-way ANOVA was used to determine significant differences. Duncan's multiple range tests (Duncan, 1955) were used to rank the treatments and mean differences were considered significant at  $p < 0.05$ .

## Results

### Protein analysis

Figure 1 shows the protein content of microalga *S. platensis* following culture with different nitrogen regimes ( $\text{NH}_4\text{Cl}$ ,  $\text{NH}_4\text{NO}_3$  and  $\text{KNO}_3$ ). The protein content was slightly higher in media containing  $\text{NH}_4\text{NO}_3$  than in that containing  $\text{NH}_4\text{Cl}$  and  $\text{KNO}_3$ . However, in all *S. platensis* cultures, increasing in nitrogen concentrations led to an increase in protein content; but there was not significant difference in terms of nitrogen regimes or concentrations ( $p > 0.05$ ).



**Figure 1:** Protein content of microalga *Spirulina platensis* following culture with different nitrogen regimes.

### Larval stage development and survival rate

Larval stage development and survival rate of *L. vannamei* fed *S. platensis* cultured with different nitrogen regimes at 5 days post hatch is shown in Table 1. The result showed that *L. vannamei* fed with different *S. platensis* cultured with different nitrogen regimes could

successfully develop to protozoa 3 ( $\text{PZ}_{\text{III}}$ ) at 5 days post hatch and in this case there was no significant difference in terms of nitrogen regimes ( $p > 0.05$ ). Moreover, no notable differences were found in terms of survival rate in different treatments.

**Table 1: Larval development and survival rate of *Litopenaeus vannamei* fed on *Spirulina platensis* cultured with different nitrogen regimes at 5 days post hatch\*.**

Treatments**	Stage composition (%)†				Survival Rate (%)
	N <sub>V</sub>	PZ <sub>I</sub>	PZ <sub>II</sub>	PZ <sub>III</sub>	
Diet A	3±0	9±1	9±1	79±5	83±7
Diet B	3±0	6±0	10±2	81±7	84±9
Diet C	3±0	8±1	9±0	80±11	82±6

\*Data are mean values of triplicate samples ± SD.; Different superscript letters in the same column indicate significant differences ( $p < 0.05$ ).

\*\* NH<sub>4</sub>Cl (Diet A); NH<sub>4</sub>NO<sub>3</sub> (Diet B); KNO<sub>3</sub> (Diet C).

†Stage composition; N<sub>V</sub>= Naplius 5; PZ<sub>I</sub> = protozoa 1, PZ<sub>II</sub> = protozoa 2 and PZ<sub>III</sub> = protozoa 3 as % of total live larvae.

The results of larval stage development and survival rate of *L. vannamei* on the 9<sup>th</sup> day after hatching that were fed on *S. platensis* cultured with different nitrogen regimes, is shown in Table 2. According to the results, shrimp larvae

fed on diets B and C developed slightly faster to PL<sub>I</sub> stage compared to larvae fed on diet A. On the other hand, there was no significant difference in terms of survival rate in different treatments at 9 days post hatch ( $p > 0.05$ ).

**Table 2: Larval stage composition and survival rate of *Litopenaeus vannamei* fed on *S. platensis* cultured with different nitrogen regimes at 9 days post hatch\*.**

Treatments	Stage composition (%)†				Survival Rate (%)
	M <sub>I</sub>	M <sub>II</sub>	M <sub>III</sub>	PL <sub>I</sub>	
Diet A	5±1	7±1	15±2	73±8	65±5
Diet B	6±0	5±0	11±1	78±6	69±7
Diet C	3±1	7±2	13±4	77±6	64±8

\*Data are mean values of triplicate samples ± SD.; Different superscript letters in the same column indicate significant differences ( $p < 0.05$ ).

\*\* NH<sub>4</sub>Cl (Diet A); NH<sub>4</sub>NO<sub>3</sub> (Diet B); KNO<sub>3</sub> (Diet C).

†Stage composition; M<sub>I</sub> = Mysis 1, M<sub>II</sub> = Mysis 2, M<sub>III</sub> = Mysis 3 and PL<sub>I</sub> = Post larvae 1 as % of total live larvae.

On the other hand, *L. vannamei* larvae that were fed on *S. platensis* cultured with NH<sub>4</sub>NO<sub>3</sub> developed significantly ( $p < 0.05$ ) faster to PL<sub>V</sub> stage compared with the other diets on the 14<sup>th</sup> day after hatching (Table 3). On the same day, survival rate of *P. vannamei* larvae was significantly higher when fed on NH<sub>4</sub>NO<sub>3</sub> compared with the other diets ( $p < 0.05$ ).

Upon termination of the experimental period (19<sup>th</sup> day after hatching), most shrimp larvae successfully molted to PL<sub>IX</sub> (77% for

Diet B, 72% for Diet C, and 70% for Diet A). A significantly higher percentage of shrimp larvae, which developed to PL<sub>IX</sub> stage and fed on *S. platensis* that was previously cultured with NH<sub>4</sub>NO<sub>3</sub> as a nitrogen regime in culture media ( $p < 0.05$ ) was found. Moreover, the survival rate was not significantly different in tested dietary regimes, following feeding on *S. Platensis* (Table 4).

**Table 3: Larval stage composition and survival rate of *Litopenaeus vannamei* fed on *S. platensis* cultured with different nitrogen regimes at 14 days post hatch\*.**

Treatments	Stage composition (%)†				Survival Rate (%)
	PL <sub>II</sub>	PL <sub>III</sub>	PL <sub>IV</sub>	PL <sub>V</sub>	
Diet A	6±2	10±2	14±2	70±9 <sup>b</sup>	48±6 <sup>b</sup>
Diet B	4±1	8±2	11±4	77±10 <sup>a</sup>	57±9 <sup>a</sup>
Diet C	4±1	11±1	13±2	72±6 <sup>b</sup>	51±5 <sup>b</sup>

\*Data are mean values of triplicate samples ± SD.; Different superscript letters in the same column indicate significant differences ( $p < 0.05$ ).

\*\* NH<sub>4</sub>Cl (Diet A); NH<sub>4</sub>NO<sub>3</sub> (Diet B); KNO<sub>3</sub> (Diet C).

†Stage composition; PL<sub>II</sub> = Post larvae 2, PL<sub>III</sub> = Post larvae 3, PL<sub>IV</sub> = Post larvae 4 and PL<sub>V</sub> = Post larvae 5 as % of total live larvae.

**Table 4: Larval stage composition and survival rate of *Litopenaeus vannamei* fed on *S. platensis* cultured with different nitrogen regimes at 19 days post hatch\*.**

Treatments	Stage composition (%)†				Survival Rate (%)
	PL <sub>VI</sub>	PL <sub>VII</sub>	PL <sub>VIII</sub>	PL <sub>IX</sub>	
Diet A	2±0	8±2	11±3	79±6 <sup>b</sup>	35±5
Diet B	1±1	6±1	8±2	85±12 <sup>a</sup>	41±2
Diet C	3±1	6±0	10±2	81±9 <sup>b</sup>	39±7

\*Data are mean values of triplicate samples ± SD.; Different superscript letters in the same column indicate significant differences ( $p < 0.05$ ).


\*\* NH<sub>4</sub>Cl (Diet A); NH<sub>4</sub>NO<sub>3</sub> (Diet B); KNO<sub>3</sub> (Diet C).

†Stage composition; PL<sub>VI</sub> = Post larvae 6, PL<sub>VII</sub> = Post larvae 7, PL<sub>VIII</sub> = Post larvae 8 and PL<sub>IX</sub> = Post larvae 9 as % of total live larvae.

### Total length

The mean final length of *L. vannamei* larvae (on the 19<sup>th</sup> day of the feeding trial) fed on *S. platensis* that were previously cultured with different nitrogen regimes is shown in figure 2. Although the maximum final length

(Diet B; 8600µ) was obtained in culture containing NH<sub>4</sub>NO<sub>3</sub>; but there was no significant differences between treatments for larval size when the nitrogen were varied in term of regimes ( $p > 0.05$ ).

**Figure 2: Final length (mm) of *Litopenaeus vannamei* larvae following culture with different nitrogen regimes.**

## Discussion

Providing a suitable nitrogen regime in culture medium can be considered as a promising method to *S. platensis* cultivation for achieving optimal biomass and protein production (Salunke *et al.*, 2016). Moreover, cultivation medium has a great impact on the productivity of biomass and other compounds of interest. For example, nitrogen concentration in the medium (optimum at 2.5 g/L) and also nitrogen source (urea better than ammonium or nitrate) has a great impact on *Spirulina* productivity (Soletto *et al.*, 2005; Çelekli and Yavuzatmaca, 2009). Therefore, in this investigation, different nitrogen regimes were chosen to evaluate those effects on the biomass and protein content of microalga *Spirulina platensis*. We determined that the protein content of *S. platensis* cultured with  $\text{NH}_4\text{NO}_3$  was slightly higher than cultures containing  $\text{NH}_4\text{Cl}$  or  $\text{KNO}_3$ . Moreover, in all *S. platensis* cultures, increasing nitrogen concentrations led to a slight increase in protein content, but it was not significantly different in terms of nitrogen regimes. The reason for the slightly higher protein amount in cultures with  $\text{NH}_4\text{NO}_3$  than cultures containing  $\text{NH}_4\text{Cl}$  or  $\text{KNO}_3$  can be explained as the  $\text{NH}_4\text{NO}_3$  contains two nitrogen atoms (35% nitrogen) than to  $\text{NH}_4\text{Cl}$  or  $\text{KNO}_3$  which have only one nitrogen atom (22 or 14% nitrogen, respectively) (Faintuch *et al.*, 1991). Moreover, the reason for the slightly lower protein amount in cultures with

low nitrogen concentrations can be interpreted as the depletion of the nitrogen in the culture medium as a result of faster growth and the prolonged steady state. Uslu *et al.* (2011) mentioned that in N-sufficient growth mediums, protein content of *S. platensis* is supported, while its content drops in N-deficient mediums.

*Spirulina* is considered as one of the most concentrated natural sources for nutrition to both terrestrial and aquatic animals. Therefore, *Spirulina* could be an excellent source of useful nutrients as well as a good energy source that can be used as a crucial component for animal feeding (Farag *et al.*, 2016). However, the main application of microalgae for aquaculture is associated with nutrition and have beneficial effects on stabilizing water quality and subsequently the health of aquatic animals (Han *et al.*, 2019). In this study, the nutritional effect of *Spirulina* cultured with different nitrogen regimes was further evaluated using *L. vannamei* larvae.

The larval development showed that *L. vannamei* fed with *S. platensis* cultured with different nitrogen regimes could successfully develop to next stages, and there was no significant difference before 9 days post hatch larvae ( $\text{PL}_1$ ). However, in general, it should be noted that *L. vannamei* larvae fed on *Spirulina* that were previously cultured with  $\text{NH}_4\text{NO}_3$  as a nitrogen regime, moulted faster to next stages compared with larvae fed on cultures containing  $\text{NH}_4\text{Cl}$  and  $\text{KNO}_3$ ,

in particular beyond the 14<sup>th</sup> day after hatching. Delayed or slightly lower larval development rates observed in cultures containing NH<sub>4</sub>Cl or KNO<sub>3</sub> could be due to inadequate nutrients or energy from *S. platensis* for *L. vannamei* larvae, which likely hindered development to the next stages. It is well known that there is a relationship between larval development rates for penaeid shrimp and protein requirement (Cuzon, 2004); and shrimp larvae have a higher dietary protein requirement than juveniles and adults (Lee and Lee, 2018). Moreover, larval development of *L. vannamei* could only be supported with a diet of *S. platensis* until the M<sub>III</sub>/PL stage, and beyond this stage, the larvae need *Artemia* nauplii as live feed. This is in agreement with the report by Ronquillo *et al.* (1997) on the larval development with other penaeid larvae of *P. semisulcatus*, *P. monodon* and *P. chinensis*.

On the same way, higher larval survival rates were obtained in *L. vannamei* larvae fed on *Spirulina* that were previously cultured with NH<sub>4</sub>NO<sub>3</sub> as a nitrogen regime. *L. vannamei* larvae fed on this diet clearly showed significant increases in survival rate beyond the 14<sup>th</sup> day after hatching, and before this stage (PL<sub>I</sub>), in general, shrimp larvae in all tested diets exhibited equal survival rates on the feeding regimes. This result indicated that the tested diets could provide nutritional requirements and subsequently promote higher survival rates for *L. vannamei* larvae. Colvin and

Brand (2009) indicated that the optimum level of dietary protein requirement by penaeid shrimp varies in different larval stages, which corresponds to the results of this study.

Indeed, similar total length of shrimp larvae was observed with the feeding trials. Jaime-Ceballos *et al.* (2005) stated that supplementation of *S. platensis* meal did not favor an increment in size of *Litopenaeus schmitti* larvae. Moreover, Teshima *et al.* (1982) indicated that growth measured as final total length may not be the best criteria for evaluating larval feeding experiments; instead, success of metamorphosis is suggested as a better indicator of the nutritional value of a larval diet. Nevertheless, the final total length of shrimp larvae observed in this study was in agreement with the larval development rate since we monitored both parameters (total length and larval development). In conclusion, this study clearly showed that using NH<sub>4</sub>NO<sub>3</sub> as a nitrogen regime in *S. platensis* culture media could provide a better diet to meet nutritional requirements and subsequently promote higher survival, growth rate and larval development for the hatchery production of *L. vannamei* larvae.

### Acknowledgments

The authors thank the Persian Gulf and Oman Sea Ecology Research Center for their technical support. The authors are grateful to Dr. Samuel Allen for his editorial assistance.



## References

- Avila-Leon, I., Matsudo, M.C., Sato, S. and Carvalho, J.C.M., 2012.** *Arthrospira platensis* biomass with high protein content cultivated in continuous process using urea as nitrogen source. *Journal of Applied Microbiology*, 112, 1086–94. DOI: 10.1111/j.1365-2672.2012.05303.x
- Bezerra, R.P., Matsudo, M.C., Converti, A., Sato, S. and Carvalho, J.C.M., 2008.** Influence of ammonium chloride feeding time and light intensity on the cultivation of *Spirulina (Arthrospira) platensis*. *Biotechnology and Bioengineering*, 100, 297–305. DOI: 10.1002/bit.21771
- Carrillo, S., Lopez, E., Casas, M.M., Avila, E., Castillo, R.M., Carranco, M.E., Calvo, C. and Perez-Gil, E., 2008.** Potential use of seaweeds in the laying hen ration to improve the quality of n-3 fatty acid enriched eggs. *Journal of Applied Phycology*, 20, 271–278. DOI:10.1007/s10811-008-9334-4
- Çelekli, A. and Yavuzatmaca, M., 2009.** Predictive modeling of biomass production by *Spirulina platensis* as function of nitrate and NaCl concentrations. *Bioresource Technology*, 100, 1847–1851. DOI:10.1016/j.biortech.2008.09.042. Epub
- Colla, L.M., Reinehr, C.O., Carolina, R. and Jorge, A.V.C., 2007.** Production of biomass and nutraceutical compounds by *Spirulina platensis* under different temperature and nitrogen regimes. *Bioresource Technology*, 98, 1489–1493. DOI: 10.1016/j.biortech.2005.09.030. Epub
- Colvin, L.B. and Brand, C.W., 2009.** The protein requirement of Penaeid shrimp at various life-cycle stages with compounded diets in controlled environment systems. *Journal of the World Aquaculture Society*, 8, 821–840. DOI:10.1111/j.1749-7345.1977.tb00164.x
- Cuzon, G., Guillaume, J. and Gaxiola, G., 2004.** Review on amino acid requirement in shrimp. (Abstract Book) *Aquaculture*, 2004. March 1-5, 2004. Hawaii Convention Center. Honolulu, Hawaii, USA. 140 P.
- Duncan, D.B., 1955.** Multiple range and multiple F tests. *Biometrics*, 11, 1–42. DOI:10.2307/3001478
- Faintuch, B.L., Sato, S. and Aquarone, E., 1991.** Influence of the nutritional sources on the growth rate of cyanobacteria. *Archives of Biology and Technology*, 34,13–30. DOI:10.1016/j.ejar.2012.09.003
- Farag, M.R., Alagawany, M., Abd El-Hack, M.E. and Dhama, K., 2016.** Nutritional and healthical aspects of *Spirulina (Arthrospira)* for poultry, animals and human. *International Journal of Pharmacology*, 12, 36–51. DOI: 10.3923/ijp.2016.36.51
- Ferreira, L.S., Rodrigues, M.S., Converti, A., Sato, S. and Carvalho, J.C.M., 2010.** A new approach to ammonium sulphate feeding for fed-batch *Arthrospira*

- (*Spirulina*) *platensis* cultivation in tubular photobioreactor. *Biotechnol. Progress*, 26, 1271–1277. DOI: 10.1002/btpr.457
- Gouveia, L., Batista, A.P., Sousa, I., Raymundo, A. and Bandarra, N.M., 2008.** Microalgae in Noval Food Products. In: Food Chemistry Research Development, Papadopoulos, K.N. (Ed.) Nova Science Publishers, Inc., Hauppauge, NY, USA. 297 P.
- Guillard R.R.L. and Ryther, J.H., 1962.** Studies on marine planktonic diatoms. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. *Canadian Journal of Microbiology* 18, 229–239. DOI: 10.1139/m62-029
- Han, P., Lu, Q., Fan L. and Zhou, W., 2019.** A Review on the Use of Microalgae for Sustainable Aquaculture. *Applied Science*, 9, 1–20. DOI:10.3390/app9112377
- Holman, B.W.B. and Malau-Aduli, A.E.O., 2012.** Spirulina as a livestock supplement and animal feed. *Journal of Animal Physiology and Animal Nutrition*, 97, 615–623. DOI: 10.1111/j.1439-0396.2012.01328.x
- Ilavarasi, A., Mubarakali, D., Praveenkumar, R., Baldev E. and Thajuddin, N., 2011.** Optimization of Various Growth Media to Freshwater Microalgae for Biomass Production, *Biotechnology*, 10, 540–545. DOI: 10.3923/biotech.2011.540.545
- Jaime-Ceballos, B., Villarreal, H., Garcia, T., Pérez –Jar, L. and Alfonso, E., 2005.** Effect of *Spirulina platensis* meal as feed additive on growth, survival and development in *Litopenaeus schmitti* shrimp larvae. *Revista de Investigaciones Marinas*, 26, 235–241.
- Jin, S.E., Lee, S.J., Kim, Y.N., Park, C.Y. 2020.** Spirulina powder as a feed supplement to enhance abalone growth, *Aquaculture Report*. 17, 100318. DOI:10.1016/j.aqrep.2020.100318
- Kermani, P., Babaei, S., Abedian-Kenari, A. and Hedayati, M., 2020.** Growth performance, plasma parameters and liver antioxidant enzymes activities of Rainbow trout (*Oncorhynchus mykiss*) juvenile fed on *Spirulina platensis* extract, *Iranian Journal of Fisheries Sciences*, 19(3), 1463-1478. DOI: 10.22092/ijfs.2019.120209
- Kulpys, J., Paulauskas, E., Pilipavicius, V. and Stankevicius, R., 2009.** Influence of cyanobacteria *Arthrospira (Spirulina) platensis* biomass additive towards the body condition of lactation cows and biochemical milk indexes. *Agronomy Research*, 7, 823–835. DOI:20.500.12259/82366
- Lavens, P. and Sorgeloos, P., 2000.** The history, present status and prospects of the availability of Artemia cysts for Aquaculture. *Aquaculture*. 181. 397-403. DOI: 10.1016/S0044-8486(99)00233-1.

- Lee, C. and Lee, K.J., 2018.** Dietary protein requirement of Pacific white shrimp *Litopenaeus vannamei* in three different growth stages. *Fisheries and Aquatic Sciences*, 21, 30. DOI:10.1186/s41240-018-0105-0
- Lowry O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J., 1951.** Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193, 265–275. DOI: 10.1016/S0021-258(19)5251-6
- Marrez, D.A., Naguib, M.M., Sultan, Y.Y., Daw Z.Y. and Higazy, A.M., 2014.** Evaluation of Chemical Composition for *Spirulina platensis* in Different Culture Media. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 5, 1161–1171.
- Matsudo, M., Bezerra, R., Sato, S., Perego, P., Converti, A. and Carvalho, J., 2009.** Repeated fed-batch cultivation of *Arthrospira (Spirulina) platensis* using urea as nitrogen source. *Biochemical Engineering Journal*, 43, 52–57. DOI:10.1016/j.bej.2008.08.009
- Mostert, E.S. and Grobbelaar, J.U., 1987.** The influence of nitrogen and phosphorus on algal growth and quality in outdoor mass algal cultures. *Biomass*, 13, 219–233. DOI:10.1016/0144-4565 (87)90061-8
- Piñero Estrada, J.E., Bescos, P.B. and Villar Del Fresno, A.M., 2001.** Antioxidant activity of different fractions of *Spirulina platensis* protean extract. *IL Farmaco*, 56, 497–500. DOI: 10.1016/s0014-827x (01)01084-9
- Rajasekaran, C., Ajeesh, C.P.M., Balaji, S., Shalini, M., Siva, R., Devanand, R.D., Fulzele, P. and Kalaivani, T. 2016.** Effect of Modified Zarrouk's Medium on Growth of Different *Spirulina*. *Journal of Science and Technology*, 13, 67–75. DOI:10.14456/VOL13IS S2PP
- Ronquillo, J.D., Matias, J.R., Saisho, T. and Yamasaki, S., 1997.** Culture of *Tetraselmis tetrahele* and its utilization in the hatchery production of different penaeid shrimps in Asia. *Hydrobiologia*, 358, 237–244. DOI: 10.1023/A:1003128701968
- Ronquillo J.D., Saisho, T. and McKinley, R.S., 2006.** Early developmental stages of the green tiger prawn, *Penaeus semisulcatus* de Haan (Crustacea, Decapoda, Penaeidae). *Hydrobiologia*, 560, 175–196. DOI: 10.1007/s10750-005-1448-y
- Salunke, K.J., Magar, S.A., Joshi, R.R. and Wadikar, M.S., 2016.** Comparative study on the growth of *Spirulina platensis* on different culture media. *Bioscience discovery*, 7, 90–92. DOI:10.1016/j.egy.2019.02.009
- Singh, S., 2006.** *Spirulina*: A Green gold mine. Paper presented at: Spirutech 2006. *Spirulina* cultivation: Potentials and Prospects. Jabalpur, Madhya Pradesh.

- Soletto, D., Binaghi, L., Lodi, A., Carvalho, J.C.M. and Converti, A., 2005.** Batch and fed-batch cultivations of *Spirulina platensis* using ammonium sulphate and urea as nitrogen sources. *Aquaculture*, 243, 217–224. DOI:10.1016/j.aquaculture.2004.10.005
- Spolaore, P., Joannis-Cassan, C., Duran, E. and Esambert, A., 2006.** Commercial applications of microalgae. *Journal of Bioscience and Bioengineering*, 101, 87–96. DOI:10.1263/jbb.101.87
- Teshima, S., Kanazawa, A., Sasada, H. and Kawasaki, M., 1982.** Requirements of the larval prawn, *Penaeus japonicus* for cholesterol and soybean phospholipids. *Memoirs of Faculty of Fisheries Kagoshima University*, 31, 193–199.
- Uslu, L., Işık, O., Koç, K. and Göksan, T., 2011.** The effects of nitrogen deficiencies on the lipid and protein of *Spirulina platensis*. *African Journal of Biotechnology*, 10, 386–389. DOI: 10.5897/AJB10.1547
- Venkataraman, L.V., Bhagyalakshmi, N. and Ravishankar, G.A., 1995.** Commercial production of micro-macro algae problems and potentials. *Indian Journal of Microbiology*, 35, 1–19. DOI: 10.15376/biores.9.1.1606-1633
- Zarrouk, C., 1966.** Contribution à l'étude d'une cyanophycée. Influence de divers facteurs physiques et chimiques sur la croissance et la photosynthèse de *Spirulina maxima*. Ph.D. Thesis, Université de Paris, Paris. 85 P