

## Research Article

# Effect of phytogenic feed additive (Digestarom PEP) on growth performance, hematological parameters, immunity system, and carcass quality of giant gourami (*Osphronemus goramy*)

Karimifar B.<sup>1</sup>; Abdolhay H.<sup>2</sup>; Mohamadizadeh F.<sup>3</sup>; Shamsaie Mehrgan M.<sup>1</sup>; Hosseini Shekarabi S.P.<sup>1</sup>

Received: July 2019

Accepted: December 2019

### Abstract

There is a growing interest among researchers to use herbal supplementations in aquafeed to enhance growth performance as well as the immunity systems of aquatic animals. Therefore, this study was conducted to investigate the effect of phytogenic feed additive (Digestarom PEP) on growth performance, chemical composition, hematological and immune responses of giant gourami (*Osphronemus goramy*). For this purpose, giant gourami with the average weight  $3.1 \pm 0.2$  g with selective experimental treatments included, control (0 g Digestarom PEP/kg), T1 PEP 0.5% (5 g Digestarom PEP/kg), T2 PEP 1% (10 g Digestarom PEP/kg), and T3 PEP 1.5% (15 g Digestarom PEP/kg) were feed for 60 days. Based on the growth performance results, a significant difference was observed in the specific growth rate (3.19 %/day), weight gain (18.47 g), and feed conversion factor (0.78) of PEP 1.5% treatment compared to other treatments ( $p < 0.05$ ). Based on the hematological results, there was a significant difference in the red and white blood cells numbers, neutrophil and monocyte percentages, and hematocrit level in PEP 1.5% treatment compared to other treatments ( $p < 0.05$ ). The levels of immune indices including lysozyme activity, IgM, complement proteins (C3 and C4) were significantly increased ( $p < 0.05$ ) in PEP 1.5% treatment compared to others, especially the control one. Also, there was a significant difference in the crude protein (15.37%) and lipid (3.87%) contents among PEP 1% and PEP 1.5% compared to control and PEP 0.5% ( $p < 0.05$ ). In conclusion, the results of this study showed that dietary administration of Digestrom™ PEP, especially at 1.5% concentration, could improve growth performance and innate immune responses of giant gourami.

**Keywords:** Digestrom PEP, Growth promoter, Immunostimulant., *Osphronemus goramy*

1- Department of Fisheries Science, Science and Research Branch, Islamic Azad University, Tehran, Iran.

2- Iranian Fisheries Sciences Research Institute (IFRSRI), Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran

3- Department of Fisheries Science, Bandarabas Branch, Islamic Azad University, Hormozgan, Iran.

\* Corresponding authors's E-mail: hossein\_abdolhay@yahoo.com

## Introduction

High stocking density, especially in the intensive aquaculture systems has generated potential environmental stress, leading to high susceptibility to various diseases including bacteria, fungi, viruses, and parasites. Recently, the application of antibiotics has been banned due to increasing the resistance to the most infectious bacteria as well as serious environmental hazards (Lazado and Caipang, 2014; Wang, 2014). For this reason, using natural feed additives as a growth promoter and immunostimulants were recommended (Syahidah *et al.*, 2015; Shekarabi *et al.*, 2022).

Phytogetic additives as an alternative of antibiotics were divided into several types, including sensory additives, technological additives (antioxidants and toxin binder), zootechnical additives (immunomodulators, digestive stimulants, growth promoters, etc.) and nutritional additives (vitamins, minerals, etc.) (Karaskova *et al.*, 2015). Improving the health status, growth performance, and immunity system of several fish species were reported in several literatures by phytogetic compounds containing carvacrol, thymol, cineole, linalool, anethole, 1,8-cineol, fennel, allicin, capsaicin, allylisothiocynate, and piperine (Grashorn, 2010; Giannenas *et al.*, 2012; Gultepe *et al.*, 2014; Sutili *et al.*, 2017; Mohammadi *et al.*, 2018; Samavat *et al.*, 2019; Shekarabi *et al.*, 2019).

Digestaron PEP is a phytogetic feed additive that contains essential oils included carvacrol, thymol, anethol, and

limonene and also a prebiotic (fructooligosaccharide). Carvacrol in this product extracted from *Mentha pulegium* has strong antimicrobial and antioxidant activities (Kamimura *et al.*, 2014). Anethol is used in Digestaron PEP. extracted from *Pimpinella anisum* has antifungal activity and appetite properties. The limonene in this product has strong antimicrobial properties and improved the taste of the food (Reisinger *et al.*, 2011). Fructooligosaccharide (FOS) is a nondigestible oligosaccharides that has positive effects on growth performance, metabolism, gastrointestinal microbiota composition and non-specific immunity of Atlantic salmon *Salmo salar* (Grisdale-Helland *et al.*, 2009) *Acipenser baerii*, beluga sturgeon (*Huso huso*) (Hoseinifar *et al.*, 2011), and *Clarias gariepinus* (Genc *et al.*, 2020). In previous studies, the positive effects of Digestaron PEP. As a feed supplement on health, immunity and chemical composition of beluga sturgeon (Defaee *et al.*, 2016), Nile tilapia (*Oreochromis niloticus*) fingerlings (Gaber *et al.*, 2014), bester juvenile fish (*H. huso* × *Acipenser ruthenus*) (Taghavi *et al.*, 2015), and channel catfish (*Ictalurus punctatus*) (Peterson and Bosworth, 2014) were reported.

Giant gourami (*Osphronemus goramy*) is an important fish species in the aquarium trade of Iran, but it is commercially farmed for food in Southeast Asia and other locations due to introductions (Roberts, 1992). The fish also has been used for controlling invasive aquatic plants like *Salvinia*

*molesta*. The growth rate of giant gourami is very fast and also it is easily adapted to formulated feed (Budi *et al.*, 2015). Therefore, the main objective of this study was to evaluate the effects of Digestarom PEP on growth performance, chemical composition, haematological and immune response of giant gourami.

## Materials and methods

### Experimental fish

One hundred and eighty healthy *O. goramy* with an average weight of  $3.1 \pm 0.2$  g were purchased from a private sector farm and transferred to Science and Research Laboratory (Tehran province, Iran). In this study, 12 aquaria ( $35 \times 50 \times 100$  cm) with air pumps for each aquarium were designed. After 14 days of acclimation, fish were distributed into 4 treatments aquaria; each contains 3 aquaria as replication (each with 15 fish). The water was maintained at  $25.0 \pm 1.1^\circ\text{C}$ , dissolved oxygen at  $7.1 \pm 0.2$  mg L<sup>-1</sup>, pH  $7.4 \pm 0.1$ .

Weight gain (WG; g) =  $W_2$  (g) -  $W_1$  (g)

Where  $W_1$  is the initial weight,  $W_2$  is the final weight (Tacon, 1990)

Specific growth rate (SGR; % d<sup>-1</sup>) =  $100 (\ln W_2 - \ln W_1) / T$

Where ln = Natural logarithm,  $W_2$  = Final weight at time  $t_2$ ,  $W_1$  = Initial weight at time  $t_1$ , and T is the number of days in the feeding period.

Feed conversion ratio (FCR) = feed intake (g) / weight gain (g)

Survival rate (SR; %) =  $100 (N_t / N_0)$

Where  $N_t$  and  $N_0$  are final and initial number of fish, respectively.

### Blood sampling

After 60 days, the fish were fasted for 24 h prior to sampling and anesthetized with clove oil (125 mg L<sup>-1</sup>) before

### Preparations of experimental diets

Digestarom PEP is a mixture of herbal feed additives prepared by Biomin GmbH Company (Herzogenburg, Austria). In this study, Digestarom PEP treatments included: control (0 g Digestarom PEP/kg), T1 PEP 0.5% (5 g Digestarom PEP/kg), T2 PEP 1% (10 g Digestarom PEP/kg), and T3 PEP 1.5% (15 g Digestarom PEP/kg) were sprayed on basal feed (21 Beyza Feed Mill Company, Iran), then dried and kept at 4°C until used. The basal diet included  $58 \pm 0.5\%$  crude protein,  $15 \pm 0.3\%$  crude lipid,  $1.6 \pm 0.1\%$  ash,  $11.5 \pm 0.4\%$  moisture and  $0.5 \pm 0.1\%$  fibre. Fish were fed on experimental diets at 4-5% of body weight and 4 times per day for 60 days. Fish were fed three times of the day.

### Growth performance

All fish were deprived of food for 24 hrs before weighing and sampling, and the following parameters were measured at the end of the feeding trial after 2 months:

sample collection (Hushangi and Hosseini Shekarabi, 2018). Blood samples (0.5 ml) were drawn from the caudal vein and were immediately

transferred to non-heparinized tubes for plasma collection. Blood samples of each replicate (9 fish per replicate, 27 per treatment) were immediately divided into two half parts. One half was transferred to a tube containing anti-coagulant (heparin) for studying the hematological analysis, while the other half was transferred to non-heparinized tubes for immunological studies. Sera samples were obtained by coagulated blood samples after centrifugation for 5 min at  $3000\times g$  at  $4^{\circ}\text{C}$  and stored at  $-80^{\circ}\text{C}$  until use (Beheshtipour, 2019).

#### *Hematological assay*

The total red blood cells (RBC,  $10^6 \text{ mm}^{-3}$ ) and white blood cells (WBC,  $10^3 \text{ mm}^{-3}$ ) were enumerated in an improved Neubauer hemocytometer using Hayem and Turck diluting fluids (Blaxhall and Daisley, 1973). Haematocrit (Ht, %) was determined by the standard microhematocrit method and expressed as percentage. The haemoglobin (Hb,  $\text{g dL}^{-1}$ ) level was determined according to the cyanomethemoglobin procedure. Furthermore, differential leukocyte cells were measured by preparing Giemsa stained smears. Blood smears were studied by light microscopy in order to make blood cell counts (Hushangi and Hosseini Shekarabi, 2018).

#### *Immunological parameters*

Lysozyme activity was measured according to the method described by Ellis (1990). Briefly, 50  $\mu\text{L}$  sera of was added to 950  $\mu\text{L}$  of a suspension of *Micrococcus lysodeikticus* ( $0.2 \text{ mg mL}^{-1}$ ) in a 0.05 M sodium phosphate buffer

(Sigma, USA) (pH 6.2) and the absorbance was measured at 520-560 nm and  $22^{\circ}\text{C}$  after 30 s and 180 s by spectrophotometer (2100-VIS model Unico, USA).

Serum immunoglobulin M (IgM) content was measured according to the method recommended by Sun *et al.* (2010) by using a microprotein determination method (C-690; Sigma), prior and after precipitating down the immunoglobulin molecules by means of a 12% solution of polyethylene glycol (Sigma). Serum levels of complements (C3 and C4) were measured using commercial kits (Pars Azmoon, Alborz, Iran) based on the immunoturbidimetry method (Rashmei *et al.*, 2020).

#### *Chemical composition*

At the end of the feeding trial, nine fish from each tank (27 fish per group) were sampled randomly. The captured fish were humanly sacrificed and stored at  $-20^{\circ}\text{C}$  prior to further analysis. The proximate composition of the fish fillets was conducted according to the standard methods described by the Association of Official Analytical Chemists (AOAC, 2005). Moisture was determined by drying the samples to a constant weight in a hot air oven (Behr, Germany) at  $105^{\circ}\text{C}$ . Crude lipid, protein and ash were determined by chloroform methanol extraction (2:1, v/v), Kjeldahl method ( $\text{N}\times 6.25$ ) and incineration in a muffle furnace at  $500^{\circ}\text{C}$  for 6 h respectively.

#### *Statistical analysis*

All data are shown as mean  $\pm$  SD. The data were subjected to statistical analysis

using the SPSS software version no. 21 (SPSS Inc., Chicago, IL, USA). The statistical analysis was done by using one-way analysis of variance (ANOVA) followed by Duncan's multiple range tests.  $P$ -value of  $<0.05$  was considered significant. All data were performed in triplicate.

## Results

### *Growth performance*

The results of growth performance and survival rate of giant gourami fed diets enriched with different levels of Digestrom™ PEP were presented in

Table 1. Based on results, weight gain (WG), specific growth rate (SGR) and feed conversion ratio (FCR) in PEP 1.5% were significantly different than other treatments ( $p<0.05$ ). According to the results, by increasing the Digestrom™ PEP concentration from 0.5% to 1.5%, the growth parameters (except for FCR) were increased ( $p<0.05$ ). Also, the highest level of survival was observed in PEP 1.5% treatment, which showed a significant difference with other treatments ( $p<0.05$ ).

**Table 1: Growth performance and survival rate of giant gourami fed diets enriched with different levels of Digestrom™ PEP after 60 days (Mean  $\pm$ SD).**

Parameter	Control	PEP 0.5%	PEP 1%	PEP 1.5%
WG (g)	7.00 $\pm$ 0.45 <sup>d</sup>	10.17 $\pm$ 0.57 <sup>c</sup>	12.33 $\pm$ 0.25 <sup>b</sup>	18.47 $\pm$ 0.85 <sup>a</sup>
SGR (%)	1.99 $\pm$ 0.18 <sup>d</sup>	2.42 $\pm$ 0.09 <sup>c</sup>	2.67 $\pm$ 0.13 <sup>b</sup>	3.19 $\pm$ 0.19 <sup>a</sup>
FCR	1.07 $\pm$ 0.04 <sup>a</sup>	0.94 $\pm$ 0.05 <sup>b</sup>	0.90 $\pm$ 0.04 <sup>b</sup>	0.78 $\pm$ 0.01 <sup>c</sup>
SR%	91.11 $\pm$ 3.85 <sup>b</sup>	93.33 $\pm$ 6.67 <sup>b</sup>	93.33 $\pm$ 6.67 <sup>b</sup>	97.78 $\pm$ 3.85 <sup>a</sup>

WG, weight gain; SGR, specific growth rate; FCR, fed conversion ratio; SR%, survival rate. Means in the same rows with different superscript are significantly different

### *Hematological profile*

The number of red and white blood cells significantly increased ( $p<0.05$ ) in Digestrom™ PEP treatments compared to control (Table 2). The highest number of white blood cells was observed in PEP 1.5% treatment which showed statistically significant differences with other treatments, especially control ( $p<0.05$ ). Based on the results, there was a significant difference in neutrophil and

monocyte percentages, and hematocrit value in PEP 1.5% treatment compared to other treatments ( $p<0.05$ ). Also, the haemoglobin concentration did not significantly change by increasing dietary levels of PEP up to 1% ( $p>0.05$ ), while the highest level was obtained in PEP 1.5% treatment ( $p<0.05$ ). No significant difference was observed in lymphocyte percentage in studied treatments ( $p>0.05$ ).

**Table 2: Hematological changes of giant gourami fed diets enriched with different levels of Digestrom™ PEP after 60 days (Mean ±SD).**

Parameter	Control	PEP 0.5%	PEP 1%	PEP 1.5%
RBC ( $10^6 \text{ mm}^{-3}$ )	5.08±0.06 <sup>b</sup>	5.43±0.05 <sup>a</sup>	5.44±0.02 <sup>a</sup>	5.42±0.01 <sup>a</sup>
WBC ( $10^3 \text{ mm}^{-3}$ )	6.44±0.03 <sup>d</sup>	9.31±0.04 <sup>c</sup>	13.1±0.10 <sup>b</sup>	13.41±0.2 <sup>a</sup>
Hct (%)	35.7±0.52 <sup>b</sup>	35.47±0.50 <sup>b</sup>	35.40±0.36 <sup>b</sup>	36.20±0.72 <sup>a</sup>
Hb (g dL <sup>-1</sup> )	5.39±0.03 <sup>b</sup>	5.83±0.03 <sup>a</sup>	5.38±0.02 <sup>b</sup>	5.35±0.06 <sup>b</sup>
Monocyte (%)	2.43±0.12 <sup>d</sup>	3.17±0.31 <sup>c</sup>	4.0±0.20 <sup>b</sup>	5.23±0.06 <sup>a</sup>
Lymphocyte (%)	77.67±2.20 <sup>a</sup>	75.67±1.53 <sup>a</sup>	76.0±1.73 <sup>a</sup>	74.67±1.15 <sup>a</sup>
Neutrophil (%)	11.56±0.12 <sup>d</sup>	12.30±0.10 <sup>c</sup>	13.33±0.21 <sup>b</sup>	15.43±0.31 <sup>a</sup>


Note: RBC, red blood cells; WBC, white blood cells; Hct, hematocrit; Hb, hemoglobin concentration. Data are presented as mean ± S.D ( $n=9$  fish from each group). Means in the same rows with different superscript are significantly different ( $p<0.05$ ).

### Immunity responses


As shown in Figures 1-4, the majority of studied immune indices were significantly increased ( $p<0.05$ ) in all treatments that received dietary Digestrom™ PEP (0.5%, 1%, and 1.5%), compared to the control one ( $p<0.05$ ). However, the highest levels of lysozyme activity, IgM, and C3 were observed in 1.5% PEP treatment ( $p<0.05$ ). The level of C4 was not significantly different among Digestrom™ PEP treatments while, it was significantly higher than the control group ( $p<0.05$ ).

### Carcass chemical composition


The chemical composition of giant gourami fed diets enriched with different levels of Digestrom™ PEP is shown in Table 2. Based on the results, there was a significant difference in the crude protein percentage and crude lipid content among PEP 1% and PEP 1.5% compared to control and PEP 0.5% ( $p<0.05$ ). Although, no significant difference was observed in moisture content and ash percentage in all studied treatments ( $p>0.05$ ) (Table 3).




**Figure 1: The effect of different concentrations of Digestrom™ PEP on lysozyme activity of giant gourami. C: control (0 g Digestarom PEP/kg), PEP 0.5% (5 g Digestarom PEP/kg), PEP 1% (10 g Digestarom PEP/kg), and PEP 1.5% (15 g Digestarom PEP/kg). Error bars indicate the standard error of the mean ( $n=3$ ). Different superscript letters indicate statistically significant differences ( $p<0.05$ ).**



**Figure 2:** The effect of different concentrations of Digestrom™ PEP on IgM level of giant gourami. C: control (0 g Digestarom PEP/kg), PEP 0.5% (5 g Digestarom PEP/kg), PEP 1% (10 g Digestarom PEP/kg), and PEP 1.5% (15 g Digestarom PEP/kg). Error bars indicate the standard error of the mean (n=3). Different superscript letters indicate statistically significant differences ( $p < 0.05$ ).



**Figure 3:** The effect of different concentrations of Digestrom™ PEP on C3 level of giant gourami. C: control (0 g Digestarom PEP/kg), PEP 0.5% (5 g Digestarom PEP/kg), PEP 1% (10 g Digestarom PEP/kg), and PEP 1.5% (15 g Digestarom PEP/kg). Error bars indicate the standard error of the mean (n=3). Different superscript letters indicate statistically significant differences ( $p < 0.05$ ).



**Figure 4:** The effect of different concentrations of Digestaron™ PEP on C4 level of giant gourami. **Figure 2:** The effect of different concentrations of Digestaron™ PEP on IgM level of giant gourami. C: control (0 g Digestaron PEP/kg), PEP 0.5% (5 g Digestaron PEP/kg), PEP 1% (10 g Digestaron PEP/kg), and PEP 1.5% (15 g Digestaron PEP/kg). Error bars indicate the standard error of the mean (n=3). Different superscript letters indicate statistically significant differences ( $p<0.05$ ).

**Table 3:** Chemical composition (% in wet basis) of giant gourami fed diets enriched with different levels of Digestaron™ PEP after 60 days (Mean  $\pm$ SD).

Parameter	Control	PEP 0.5%	PEP 1%	PEP 1.5%
Crude protein	13.6 $\pm$ 0.45 <sup>c</sup>	14.27 $\pm$ 0.12 <sup>bc</sup>	15.1 $\pm$ 0.20 <sup>a</sup>	15.37 $\pm$ 0.74 <sup>a</sup>
Crude lipid	4.7 $\pm$ 0.23 <sup>a</sup>	4.27 $\pm$ 0.12 <sup>ab</sup>	4.03 $\pm$ 0.21 <sup>b</sup>	3.87 $\pm$ 0.21 <sup>b</sup>
Ash	4.87 $\pm$ 0.12 <sup>a</sup>	4.73 $\pm$ 0.06 <sup>a</sup>	4.77 $\pm$ 0.15 <sup>a</sup>	4.77 $\pm$ 0.12 <sup>a</sup>
Moisture	76.3 $\pm$ 2.0 <sup>a</sup>	76.7 $\pm$ 1.53 <sup>a</sup>	76.1 $\pm$ 1.73 <sup>a</sup>	76.0 $\pm$ 1.0 <sup>a</sup>

Means in the same rows with different superscript are significantly different ( $p<0.05$ ).

## Discussion

Phytogenics can act as natural growth promoters that improve diet palatability, stimulate appetite and increase feed consumption and growth performance (Karaskova *et al.*, 2015). According to the results, the growth indices were increased by increasing the Digestaron™ PEP concentration from 0.5% to 1.5%, while FCR were decreased by increasing the Digestaron™ PEP concentration.. At the end of the study, 1.5% PEP treatment has highest levels of weight gain, specific growth factor and survival rate. These results may be related to stimulation of digestive secretions, increasing the population of positive bacteria such as LAB, improvement of

feed digestibility, especially for proteins and amino acid (Peterson *et al.*, 2015; Ran *et al.*, 2016; Sutuli *et al.*, 2017). In a same study, El-Hawarry *et al.* (2018) showed that phytogenic compounds including oregano oil in Nile tilapia diet significantly improved the growth performance. Similar to our results, Gaber *et al.* (2014) showed that final weight and SGR of Nile tilapia were significantly affected in treatment containing 30% level wet date and 0.03% Digestaron™ PEP. Also, in Peterson and Bosworth (2016) study, channel catfish fed diets enriched with Digestaron™ significantly increased the weight gain and decreased the FCR. Also, Defaee *et al.* (2016) reported that



the Digestrom™ PEP at 2% of juvenile beluga diet could enhance growth parameters (average daily growth, specific growth rate and protein efficiency) and survival of the juvenile fish. Results of the present study were similar to Zheng *et al.* (2009) and Giannenas *et al.* (2012) studies that added carvacrol and thymol in catfish (Zheng *et al.*, 2009) and rainbow trout (*Oncorhynchus mykiss*) feed, respectively (Giannenas *et al.*, 2012). Present results showed that Digestrom™ PEP increased the survival rate of giant gourami. The highest level of survival rate was observed in the PEP 1.5% treatment, which had a significant difference from the control. It was in agreement with Peterson *et al.* (2015) and Giannenas *et al.* (2012) studies that added Digestrom™ PEP in channel catfish and Biomin™ (commercial products containing carvacrol and thymol) in rainbow trout feed, respectively.

Hematology indicators are known as valuable biological indicators that reflect the body's general health (Haghighi *et al.*, 2017). Based on the results, there was a significant difference in neutrophil and monocyte percentage, haematocrit and haemoglobin values in PEP 1.5% treatment compared to other treatments ( $p < 0.05$ ). Increasing the number of white blood cells and neutrophils in this study may be related to the presence of fructooligosaccharide, carvacrol and thymol in the PEP supplement that has stimulation effects on the immune system. In Hoseinifar *et al.* (2011) study, Hct, Hb and WBC were significantly

higher in fish fed 2% oligofructose than the control group ( $p < 0.05$ ). Defaee *et al.* (2016) reported that WBC and neutrophils of juvenile beluga in the PEP 1% and 2% treatment was significantly higher than other treatments. Also, Gultepe *et al.* (2014) found higher counts of white and red blood cells and hematocrit levels in the tilapia (*Oreochromis mossambicus*) groups that received the 1% extract of thyme (*Thymus vulgaris*), rosemary (*Rosmarinus officinalis*), and fenugreek (*Trigonella foenum graecum*) (Beheshtipour *et al.*, 2018). In a similar study, the erythrocyte number, hemoglobin content, hematocrit and leukocyte numbers were increased significantly after adding garlic to fish diets (Martins *et al.*, 2002) which could be due to the presence of limonene in both garlic and Digestrom™ PEP (Peterson and Bosworth, 2016; Utama-Ang *et al.*, 2020).

Lysozyme is responsible for activating other important defense molecules, including the complement system and phagocytic cells (Ellis, 1990; Shekarabi *et al.*, 2021). At the end of the study, lysozyme activity, IgM, C3 and C4 levels were increased in all treatments that received Digestrom™ PEP (0.5%, 1% and 1.5%), compared to control group ( $p < 0.05$ ). These results are mainly attributed to carvacrol and thymol in Digestrom™ PEP (Volpatti *et al.*, 2013). In the same study, lysozyme activity and complement levels were higher in phytogenic supplemented groups containing carvacrol and thymol compared to the control (Giannenas *et*

*al.*, 2012). Also, a significant increase in the phagocytic and lysozyme activities were observed in rainbow trout that received allicin (Nya *et al.*, 2010), Persian shallot powder (*Allium stipitatum*; Shekarabi *et al.*, 2022), and dandelion flower extract (*Taraxacum officinale*; Shekarabi *et al.*, 2021) compared to the control. Baba *et al.* (2016) showed that limon peel essential oil supplementation increased the Nitro blue tetrazolium (NBT) positive cell count and lysozyme activity of tilapia compared to control. Haghghi *et al.* (2018) reported that dietary inclusion of *O. vulgare* extract 1% improved the nonspecific immune parameters (respiratory burst activity, phagocytic activity and serum lysozyme activity) of juvenile rainbow trout. Also, Peterson *et al.* (2018) demonstrated that different herbal essential oils in Digestrom™ PEP could enhance survival of channel catfish challenged with *Edwardsiella ictaluri* through upregulation of RBL1a and RBL3b in the gill.

Based on the results, there was a significant difference in the crude protein percentage and crude lipid content among PEP 1% and PEP 1.5% compared to control and PEP 0.5% ( $p < 0.05$ ). In Peterson *et al.* (2015) survey, chemical analysis of channel catfish showed that fillet lipid was lower and protein content was higher than the control group. In the same study, a combination of 30% level wet date and 0.03% Digestrom™ PEP in Nile tilapia feed significantly increased the protein content, the protein efficiency ratio (PER) and protein productive value

(PPV%) compared to the control group (Gaber *et al.*, 2014). Also, the beluga chemical analysis revealed a significantly higher protein content in the PEP 2% treatment than control group (Defaee *et al.*, 2016). It was confirmed that phytogetic feed additive induce transcription, increasing RNA, leading to an increase in total amino acids and enhancing the protein content (Karaskova *et al.*, 2015).

In conclusion, the results suggested that Digestrom™ PEP supplement, especially in the 1.5% concentration, has positive effects on growth performance, chemical composition and immune response of giant gourami (*O. goramy*). However, further studies are needed to determine the non-specific mechanism of Digestrom™ PEP on giant gourami and survey the effects of PEP on fish physiology.

### Acknowledgements

This study was conducted in Science and Research Branch Laboratory, Islamic Azad University (Tehran, Iran). Therefore, a special thank goes to the laboratory staffs.

### References

- AOAC, 2005.** Official Methods of Analysis. 17th ed. Association of Official Analytical Chemistry, Arlington, Virginia, USA, 2200 P.
- Baba, E., Acar, Ü., Öntaş, C., Kesbiç Osman S. and Yılmaz, S., 2016.** Evaluation of Citrus limon peels essential oil on growth performance, immune response of Mozambique tilapia *Oreochromis mossambicus*

- challenged with *Edwardsiella tarda*. *Aquaculture*, 465, 13-18. DOI: 10.1016/j.aquaculture.2016.08.023
- Beheshtipour, J., Akradi, L. and Raeeszadeh, M., 2018.** The effect of aqueous extract of grapevine leaf (*Vitis vinifera*) on pathologic feature of the pancreas in type-1 experimental diabetes: a different approach to medicinal plants. *Scientific Journal of Kurdistan University of Medical Sciences*, 23(4), 91-99. DOI: 10.52547/sjku.23.4.91
- Beheshtipour, J., 2019.** Appropriate approach to technology in animal hematology. *Toxicology Reports*, 6, 118. DOI: 10.1016/j.toxrep.2018.12.008
- Blaxhall, P.C and Daisley, K.W., 1973.** Routine hematological methods for use with fish blood. *Journal of Fish Biology*, 5, 771-781.
- Budi, D.S., Muddin, A. and Suprayudi, M.A., 2015.** Growth response and feed utilization of giant gourami (*Osphronemus goramy*) juvenile feeding different protein levels of the diets supplemented with recombinant growth hormone. *Hayati Journal of Biosciences*, 22(1), 12-19. DOI: 10.4308/hjb.22.1.12
- Defaee, S., Falahatkar, B. and Efatpanah, I., 2016.** Effects of digestrom PEP on growth and some hematological parameters of juveniles' Beluga sturgeon (*Huso huso*). *Fisheries Sciences and Technology*, 5(1), 83-95. (In Farsi)
- El-Hawarry, W. N., Mohamed, R. A. and Ibrahim, S. A., 2018.** Collaborating effects of rearing density and oregano oil supplementation on growth, behavioral and stress response of Nile tilapia (*Oreochromis niloticus*). *The Egyptian Journal of Aquatic Research*, 44(2), 173-178. DOI: 10.1016/j.ejar.2018.06.008
- Ellis, A.E., 1990.** Lysozyme assay in techniques in fish immunology. 2th ed. In: Stolen JS, Fletcher TC, Anderson DP, Robertson BS, Van Muiswinkel WR, (Eds.). Fair Haven, NJ, USA: SOS Publications, 103 P.
- Gaber, M.M., Labib, E.H., Omar, E.A., Zaki, M.A. and Nour. A.M., 2014.** Effect of partially replacing corn meal by wet date on growth performance in Nile tilapia (*Oreochromis niloticus*) fingerlings, diets supplemented with Digestarom®. *Journal of Geoscience and Environment Protection*, 2, 60-67. DOI: 10.4236/gep.2014.22010
- Genc, E., Genc, M.A., Kaya, D., Secer, F.S., Qaranjiki, A. and Guroy, D., 2020.** Effect of prebiotics on the growth performance, haematological, biochemical, and histological parameters of African catfish (*Clarias gariepinus*) in recirculating aquaculture system. *Turkish Journal of Veterinary and Animal Sciences*, 44, 1222-1231. DOI:10.3906/vet-2005-106
- Giannenas, I., Triantafillou, El., Stavrakakis, S., Margaroni, M., Mavridis, S., Steiner T. and Karagouni E., 2012.** Assessment of dietary supplementation with carvacrol or thymol containing feed

- additives on performance, intestinal microbiota and antioxidant status of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 350–353, 26–32. DOI: 10.1016/j.aquaculture.2012.04.027
- Grashorn, M.A., 2010.** Use of phytobiotics in broiler nutrition – an alternative to in feed antibiotics? *Journal of Animal Feed Sciences*, 19, 338-347. DOI: 10.4236/oalib.1103640
- Grisdale-Helland, B., Helland, S.J. and Gatlin, D.M., 2009.** The effects of dietary supplementation with mannan oligosaccharide, fructo-oligosaccharide or galacto oligosaccharide on the growth and feed utilization of Atlantic salmon *Salmo salar*. *Aquaculture*, 283, 163-167. DOI: 10.1016/j.aquaculture.2008.07.012
- Gultepe, N., Bilen, S., Yilmaz, S., Guroy, D. and Aydin, S., 2014.** Effects of herbs and spice on health status of tilapia (*Oreochromis mossambicus*) challenged with *Streptococcus iniae*. *Acta Veterinaria Brno*, 83, 125–131. DOI: 10.2754/avb201483020125
- Haghighi, M., Sharif Rohani, M., Pourmoghim, H., Samadi, M., Tavoli, M., Eslami, M. and Yusefi, R., 2017.** Enhancement of immune responses of rainbow trout (*Oncorhynchus mykiss*) fed a diet supplemented with *Aloe vera* extract. *Iranian Journal of Fisheries Sciences*, 16 (3), 884-896. DOI: 10.22092/IJFS.2018.114706
- Haghighi, M., Pourmoghim, H. and Sharif Rohani, M., 2018.** Effect of *Origanum vulgare* extract on immune responses and hematological parameters of rainbow trout (*Oncorhynchus mykiss*). *Oceanography and Fisheries Open access journal*, 6, 555-687. DOI: 10.19080/OFOAJ.2018.06.555687
- Hoseinifar, S.H., Mirvaghefi, A., Merrifield, D.L., Mojazi Amiri, B., Yelghi, S. and Darvish Bastami, K., 2011.** The study of some haematological and serum biochemical parameters of juvenile beluga (*Huso huso*) fed oligofructose. *Fish Physiology and Biochemistry*, 37, 91-96. DOI: 10.1007/s10695-010-9420-9
- Hushangi, R. and Hosseini Shekarabi, S. 2018.** Effect of a peracetic acid-based disinfectant on growth, hematology and histology of juvenile rainbow trout (*Oncorhynchus mykiss*). *Fishes*, 3(1), 10. DOI: 10.3390/fishes3010010
- Kamimura, J.A., Santos, E.H., Hill, L.E. and Gomes, Carmen.L., 2014.** Antimicrobial and antioxidant activities of carvacrol microencapsulated in hydroxypropyl-beta-cyclodextrin. *LWT - Food Science and Technology*, 57(2), 701-709. DOI: 10.1016/j.lwt.2014.02.014
- Karaskova, K., Suchy, P. and Strakova, E., 2015.** Current use of phytogenic feed additives in animal nutrition: a review. *Czech Journal of Animal Science*, 60 (12), 521–530. DOI: 10.17221/8594-CJAS

- Lazado, C.C. and Caipang, C.M.A., 2014.** Atlantic cod in the dynamic probiotics research in aquaculture. *Aquaculture*, 424–425, 53-62. DOI: 10.1016/j.aquaculture.2013.12.040
- Martins, M.L., Moraes, F.R., Miyazaki, D.M., Brum, C.D., Onaka, E.M. and Fenerick, J., 2002.** Alternative treatment for *Anacanthorus penilabiatu* (*Monogenea: Dactylogyridae*) infection in cultivated pacu, *Piaractus mesopotamicus* (*Osteichthyes: Characidae*) in Brazil and its haematological effects. *Parasitology*, 9, 175-180.
- Mohammadi, M., Soltani, M., Siahpoosh, A., Hosseini Shekarabi, S. P., Shamsaie Mehrgan, M. and Lymbery, A. 2018.** Effect of date palm (*Phoenix dactylifera*) seed extract as a dietary supplementation on growth performance immunological haematological biochemical parameters of common carp. *Aquaculture Research*, 49(8), 2903-2912. DOI: 22092/IJFS.2018.115477
- Nya, E.J., Dawood, Z. and Austin, B., 2010.** The garlic component, allicin, prevents disease caused by *Aeromonas hydrophila* in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 33(4), 293-300. DOI: 10.1111/j.1365-2761.2009.01121.x
- Peterson, B.C., Peatman, E., Ourth, D.D. and Waldbieser, G.C., 2015.** Effects of a phyto-genic feed additive on growth performance, susceptibility of channel catfish to *Edwardsiella ictaluri* and levels of mannose binding lectin. *Fish & Shellfish Immunology*, 44, 21–25. DOI: 10.1016/j.fsi.2015.01.027
- Peterson, B. and Bosworth, B., 2014.** Assessment of a phyto-genic feed additive (Digestarom PEP. MGE) on growth performance, processing yield, fillet composition, and survival of channel catfish. *Journal of the World Aquaculture Society*, 45(2), 206-212. DOI: 10.1111/jwas.12103
- Peterson, B.C., Peatman, E., Ourth, D.D. and Waldbieser, G.C., 2018.** Phyto-genic feed-additive effects on channel catfish rhamnose-binding lectin levels, and susceptibility to *Edwardsiella ictaluri*. *Diseases of Aquatic Organisms*, 129(2), 99-106. DOI: 10.3354/dao03235
- Ran, C., Hu, J., Liu, W., Liu, Z., He, S. and Dan, B.C.T., 2016.** Thymol and carvacrol affect hybrid tilapia through the combination of direct stimulation and an intestinal microbiota-mediated effect: insights from a germ-free zebrafish model. *The Journal of Nutrition*, 146, 1132–1140. DOI: 10.3945/jn.115.229377
- Rashmei, M., Shekarabi, S. P. H., Mehrgan, M. S. and Paknejad, H., 2020.** Stimulatory effect of dietary chasteberry (*Vitex agnus-castus*) extract on immunity, some immune-related gene expression, and resistance against *Aeromonas hydrophila* infection in goldfish (*Carassius auratus*). *Fish & Shellfish Immunology*, 107, 129-136. DOI: 10.1016/j.fsi.2020.09.037

- Reisinger, N., Steiner, T., Nitsch, S., Schatzmayr, G. and Applegate, T.J., 2011.** Effects of a blend of essential oils on broiler performance and intestinal morphology during coccidial vaccine exposure. *Poultry Science Association*, 20, 272-283. DOI: 10.3382/japr.2010-00226
- Roberts, T.R., 1992.** Systematic revision of the Southeast Asian anabantoid fish genus *Osphronemus*, with descriptions of two new species. *Ichthyological Exploration of Freshwaters*, 2, 351-360.
- Samavat, Z., Shamsaie Mehrgan, M., Jamili, S., Soltani, M. and Hosseini Shekarabi, S.P., 2019.** Determination of grapefruit (*Citrus paradisi*) peel extract bio-active substances and its application in Caspian white fish (*Rutilus frisii kutum*) diet: Growth, haemato-biochemical parameters and intestinal morphology. *Aquaculture Research*, 50(9), 2496-2504. DOI: 10.1111/are.14204
- Shekarabi, S.P.H., Omid, A.H., Dawood, M.A., Adel, M., Avazeh, A. and Heidari, F., 2019.** Effect of black mulberry (*Morus nigra*) powder on growth performance, biochemical parameters, blood carotenoid concentration, and fillet color of rainbow trout. *Annals of Animal Science*, 20(1), 125-136. DOI: 10.2478/aoas-2019-0068
- Shekarabi, S. P. H., Mostafavi, Z. S., Mehrgan, M. S. and Islami, H. R., 2021.** Dietary supplementation with dandelion (*Taraxacum officinale*) flower extract provides immunostimulation and resistance against *Streptococcus iniae* infection in rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology*, 118, 180-187. DOI: 10.1016/j.fsi.2021.09.004
- Shekarabi, S. P. H., Javarsiani, L., Mehrgan, M. S., Dawood, M. A. and Adel, M., 2022.** Growth performance, blood biochemistry profile, and immune response of rainbow trout (*Oncorhynchus mykiss*) fed dietary Persian shallot (*Allium stipitatum*) powder. *Aquaculture*, 548, 737627. DOI: 10.1016/j.aquaculture.2021.737627
- Sun, Y.Z., Yang, H.L., Ma, R.L. and Lin, W.Y., 2010.** Probiotic applications of two dominant gut *Bacillus* strains with antagonistic activity improved the growth performance and immune responses of grouper *Epinephelus coioides*. *Fish & shellfish immunology*, 29(5), 803-809. DOI: 10.1016/j.fsi.2010.07.018
- Sutili, F.J., Delbert, M., Gatlin, III., Berta, M., and Bernardo Baldisserotto, H., 2017.** Plant essential oils as fish diet additives: benefits on fish health and stability in feed. *Reviews in Aquaculture*, 10(3), 716-726. DOI: 10.1111/raq.12197
- Syahidah, A., Saad, C.R., Daud, H.M. and Abdelhadi, Y.M., 2015.** Status and potential of herbal applications in aquaculture: A review. *Iranian Journal of Fisheries Sciences*, 14(1), 27-44. DOI: 10.22092/IJFS.2018.114421
- Tacon, A.G., 1990.** Standard methods for the nutrition and feeding of

- farmed fish and shrimp. Feeding methods. 2 th ed. Agent Laboratories Press, Redmond, Taoka, P123.
- Taghavi, F., Nasri, M. and Zamini, A.A., 2015.** Determination on growth indices, blood parameters, Immune responses and intestine bacterial flora of Bester juvenile fish (*Huso huso*×*Acipenser ruthenus*) fed by Biomin nutritional supplements. *Journal of Advances in Bioresearch*, 6 (3), 300-312. DOI: 10.15515/abr.0976-4585.6.3.136143
- Utama-ang, N., Cheewinworasak, T., Simawonthamgul, N. and Samakradhamrongthai., R.S., 2020.** Influence of garlic and pepper powder on physicochemical and sensory qualities of flavoured rice noodle. *Scientific Reports*, 10(8538), 1-9. DOI: 10.1038/s41598-020-65198-4
- Volpatti, D., Chiara, B., Francesca, T. and Marco, G., 2013.** Growth parameters, innate immune response and resistance to *Listonella* (*Vibrio*) *anguillarum* of *Dicentrarchus labrax* fed carvacrol supplemented diets. *Aquaculture Research*, 45, 31–44. DOI: 10.1111/j.1365-2109.2012.03202.x
- Wang, F.H., 2014.** High throughput profiling of antibiotic resistance genes in urban park soils with reclaimed water irrigation. *Environmental Sciences and Technology*, 48, 9079–9085. <https://doi.org/10.1021/es502615e>
- Zheng, Z.L., Justin, Y.W., Tan, H.Y., Liu, X.H., Zhou, X.X. and Wang, K.Y., 2009.** Evaluation of oregano essential oil (*Origanum heracleoticum* L.) on growth, antioxidant effect and resistance against *Aeromonas hydrophila* in channel catfish (*Ictalurus punctatus*). *Aquaculture*, 292, 214–218. DOI: 10.1016/j.aquaculture.2009.04.025