

## Research Article



# Preparation and biodistribution study of $^{67}\text{Ga}$ -rosemary (*Salvia rosmarinus* Spenn.) extract nanoparticles and its SPECT imaging in healthy rainbow trout (*Oncorhynchus mykiss*)

Damavandi Kamali N.<sup>1</sup>; Rajabifar S.<sup>2\*</sup>; Alishahi A.R.<sup>1</sup>; Heidarieh M.<sup>2\*</sup>; Ojagh S.M.<sup>1</sup>; Mirsadeghi S.H.<sup>1</sup>

Received: June 2022

Accepted: August 2022

### Abstract

The aim of this study was to evaluate the labeling of rosemary extract nanoparticles (RE-NPs) using  $^{67}\text{Ga}$  radioisotope, and their biodistribution in healthy rainbow trout (*Oncorhynchus mykiss*) tissues. RE-NPs were prepared by gamma irradiation (irradiated at a dose of 30 kGy of gamma-ray by  $^{60}\text{Co}$  (PX-30 IssIedovapel, Russia, dose rate of 0.02 Gy sec<sup>-1</sup>) and ultra-sonication methods at two concentrations of 0.5% and 1% and labeled by  $^{67}\text{Ga}$  radioisotope. Quality control studies were done using the RTLC method. The radiolabeling RE-NPs at 1% concentration showed 97% efficiency using a mixture of sodium acetate and acetic acid buffer at pH=6.5 after 30 min at room temperature. In the biodistribution study, 54 healthy fish were randomly distributed into two treatment groups (n=18 fish/treatment group) and one group served as healthy control. Fish received 3.7-7.4 MBq (300  $\mu\text{L}$ ) of radiolabeled RE-NPs at 1% concentration and free  $^{67}\text{GaCl}_3$ , intraperitoneally. The  $^{67}\text{Ga}$ -RE-NPs single-photon emission computed tomography (SPECT) demonstrated the highest tracer accumulation in the kidney and the least uptake in the brain at 24 h after injection of  $^{67}\text{Ga}$ -RE-NPs. However, 48 h after injection of  $^{67}\text{Ga}$ -RE-NPs, the uptake was negligible in all organs as compared to free  $^{67}\text{GaCl}_3$ . The obtained results revealed that administration of  $^{67}\text{Ga}$ -RE-NPs, as a natural tracer, by IP injection can be a good imaging method to visualize and understand the whole-body, especially the kidney and spleen tissues distribution and pharmacokinetics in healthy rainbow trout.

**Keywords:** Rosemary,  $^{67}\text{Ga}$ , radiolabeling, Biodistribution studies, Rainbow trout, Gamma irradiation, SPECT

1-Department of Seafood Science and Technology, Faculty of Fisheries and Environmental Sciences, Gorgan University of Agricultural Sciences and Natural Resource, Gorgan, Iran.

2-Nuclear Science and Technology Research Institute, Tehran, Iran.

\*Corresponding author's Email: mheidarieh@aeoi.org.ir; srajabifar@aeoi.org.ir

## Introduction

Antioxidants are extensively used to prevent the negative consequences of lipid oxidation (Shahidi, 1997). Natural antioxidants are being extensively studied, and spices and herbs are identified as important sources of these compounds. Diverse studies evaluated the antioxidant properties of rosemary, *Salvia rosmarinus* Spenn. (Madsen and Bertelsen, 1995; Xie *et al.*, 2017; Rezanejad *et al.*, 2019a). Rosemary contains certain compounds with high antioxidant activity like carnosic acid and carnosol (Offord *et al.*, 1997; Tironi *et al.*, 2010). Rosemary is the spice with the highest antioxidant activity and a member of the Lamiaceae family which is an attractive evergreen shrub with pine needle-like leaves that grows wild in most Mediterranean countries (Pintore *et al.*, 2002). Radiolabeling is a process that is frequently used in medicine, drug research and development and environmental case studies (Saha, 2004). Additionally, isotopic labeling is a technique used to track the passage of an isotope through a reaction in metabolic pathways or cells (Kitson *et al.*, 2014). Among them, gallium 67 ( $^{67}\text{Ga}$ ) is increasingly utilized in a great number of fields. The 78.3 h physical half-life and a rather good detectability of its photon emission make gallium 67 one of the most suitable nuclides for radiopharmaceutical research (Heidarieh *et al.*, 2014). The focus of the present work was to study the labeling of gamma-irradiated (at 30 kGy) rosemary extract nanoparticles (RE-NPs) using  $^{67}\text{Ga}$  radioisotope, and their

biodistribution in healthy rainbow trout (*Oncorhynchus mykiss*) tissues.

## Materials and methods

### *Fish*

54 healthy rainbow trout (*Oncorhynchus mykiss*) weighing 400–450 g were raised from a commercial fish farm in Karaj, Iran, and kept in running water (flow rate 0.4 l/s) in polypropylene tanks (300 L) with a water temperature of  $15\pm 2^\circ\text{C}$ , dissolved oxygen of 8.2 ppm, and natural photoperiod of 10 L:14 D.

### *Preparation of gamma-irradiated rosemary extract nanoparticles (RE-NPs)*

Rosemary (*Salvia rosmarinus* Spenn) extract nanoparticles were prepared using ultrasonic and gamma irradiation methods which were described by Rezanejad *et al.* (2019a). Rosemary powder was suspended in phosphate-buffered saline (PBS) (0.2 M, sterile, pH=7.2). Samples were sonicated for 30 min ultrasonic water bath (Jencons, England) and centrifuged at 6000 rpm for 20 min. Subsequently, precipitated in 98% ethanol, ground rosemary powder was dried at  $45^\circ\text{C}$  and milled to the micromesh sieves in aperture sizes from 53 to 96  $\mu\text{m}$ . Then, Rosemary extract powder was irradiated at a dose of 30 kGy by  $^{60}\text{Co}$  (PX-30 IssIedovapel, Russia, dose rate of  $0.02\text{ Gy sec}^{-1}$ ). RE-NPs solutions of 0.5% and 1% were prepared in PBS buffer (0.1 M, pH=6.5) (Rezanejad *et al.* 2019a).

### *Transmission electron microscopy*

A transmission electron microscope (TEM) was used to analyze the morphology and size of RE-NPs (Amelinckx and Van Landuyt, 2003). The nanoparticles were immobilized on a coated copper grid and were allowed to dry at room temperature. The particle size and shape were observed using an FEI/Philips EM 208S TEM.

### *Labelling of RE-NPs using $^{67}\text{GaCl}_3$*

For the radiolabeling experiment,  $^{67}\text{Ga}$  was received from a cyclotron, in Karaj, Iran (Jalilian *et al.*, 2007). RE-NPs were labeled using an optimization protocol according to Orlando *et al.* (1994), with minor modifications. A fraction containing  $^{67}\text{GaCl}_3$  (37-110 MBq) in a volume of 100  $\mu\text{L}$  was dried by air flow and mild heat was used. Radiolabeling was performed by mixing 100  $\mu\text{l}$   $^{67}\text{GaCl}_3$  and 1 mL of phosphate buffer and 50  $\mu\text{L}$  of RE-NPs and incubating at 90°C for 30 min. The resulting solution was stirred at room temperature for 30 min.

### *Quality control of $^{67}\text{Ga-RE-NPs}$*

The radiolabeling yield was determined by thin layer chromatography analysis (TLC), for the purity of the radio-labeled sample, one small drop of radiopharmaceutical mixture of RE-NPs labeled with  $^{67}\text{GaCl}_3$  (0.5% and 1%) was dropped on the lower side of chromatography paper strips (Whatman No. 1. Whatman, Maidstone, UK). A mixture of sodium acetate and acetic acid buffer (1 M) was used as a mobile phase which moved up and passed through the droplet and carried on free

$^{67}\text{GaCl}_3$ . The strip was dried after the mobile liquid reached the top and it was immediately scanned using RTLC (Bioscan AR-2000).  $^{67}\text{Ga-RE-NPs}$  remained at the application point, while unbound  $^{67}\text{Ga}$  ions migrated with the solvent front. The labeling efficiency of  $^{67}\text{Ga-RE-NPs}$  was determined from the upper and lower-end counts. A control test was also carried out, in the absence of RE-NPs (Jalilian *et al.*, 2007).

### *Ex vivo biodistribution studies*

All studies were approved by the University of Tehran, Biomedical Research Ethics Committee (IR.UT.VETMED.REC.). The 45 healthy rainbow trout specimens were acclimated for 7 days and randomly divided into two treatment groups and control (n=18 fish/each group). Each fish was IP injected with 300  $\mu\text{l}$  free  $^{67}\text{GaCl}_3$  and the radiolabeled nanoparticles. The total amount of radioactivity injected into each rainbow trout was measured by a 1-ml syringe before and after injection in a dose calibrator (CRC-15R, Capintec Company) with fixed geometry.

### *SPECT imaging of $^{67}\text{Ga-RE-NPs}$ in rainbow trout*

The biodynamic routes of the  $^{67}\text{Ga-RE-NPs}$  inside the fish were studied using a SPECT camera. The rainbow trout was selected randomly and transferred to a small plastic aquarium and euthanized with an overdose of MS222 (200 mg/L) (Sigma-Aldrich, Denmark) at specified times and the radioactivity was quantitated in tissues. Fish were scanned

by a dual-head gamma camera system (model: DST-XL made by SMV company). SPECT images of the whole-body were obtained 24 h and 48 h after injections of free  $^{67}\text{GaCl}_3$  and  $^{67}\text{Ga}$ -RE-NPs. The Medium-Energy All Purpose (MEAP) collimator was used at the head of the static planar acquisition with no gantry movement.

#### *Biodistribution of $^{67}\text{Ga}$ - RE-NPs in rainbow trout*

Anatomical biodistribution of  $^{67}\text{Ga}$ - RE-NPs was determined by administering nanoparticles to healthy rainbow trout. For preventing any aggregation of these nanoparticles, the labeled nanoparticles were sonicated before IP injecting (Boyer *et al.*, 2010). At times of 24 h and 48 h, the fishes were injected with rosemary extract nanoparticles and were euthanized and the major organs were collected (3 fish/time/tank). Then, the extracted organs (liver, kidney, heart, spleen, intestine, skin, muscle, stomach, brain and gill) were washed with normal saline, dried on filter paper, weighed, and their radioactivity of them measured using High Purity Germanium (HPGe) radiation detectors (Canberra, GC1020-7500SL). Accumulation of  $^{67}\text{Ga}$ -RE-NPs in each sample was expressed as the percentage injected activity per gram of tissue (% ID/g $\pm$ Standard Deviation) and calculated compared to activities of the standard dose of the injected solution (Jalilian *et al.*, 2011).

#### **Results**

The TEM images, as shown in Figure 1, reveal the morphology of RE-NPs. The

average diameter of the particles was about 55-70 nm. The best labeling results were obtained at RE-NPs solution (1% concentration, irradiated at a dose of 30 kGy of gamma ray) after 30 min at pH=6.5 (Fig. 2). The measured labeling efficiency of RE-NPs labeled with  $^{67}\text{Ga}$  was ~97%, as determined by RTLC method, which could be used without further purification for later studies. Rainbow trout showed uptake of free  $^{67}\text{GaCl}_3$  and  $^{67}\text{Ga}$ -RE-NPs in the skin, muscle, heart, brain, gill, stomach, spleen, liver and kidney at 24 h and 48 h following IP injection (Fig. 3). The organ with the highest concentration of  $^{67}\text{Ga}$ -RE-NPs was kidney ( $10.12\pm 3.21$  %ID/g), followed by spleen ( $7.83\pm 1.32$  %ID/g), liver ( $4.32\pm 0.54$  %ID/g), and muscle ( $3.91\pm 0.26$  %ID/g) at 24 h. There was low-level tracer uptake into the brain ( $0.11\pm 0.02$  %ID/g) and heart ( $0.83\pm 0.01$  %ID/g) with minimal uptake into the other organs at 24 h. After 48 h, the accumulation level of  $^{67}\text{Ga}$ -RE-NPs was negligible for all of the organs except the liver and skin. Specifically, accumulation of  $^{67}\text{Ga}$ -RE-NPs in the liver was increased with time ( $4.32\pm 0.54$  % ID/g at 24 h, and  $6.71\pm 0.23$  % ID/g at 48 h, after injections). On the contrary, in the heart, stomach, spleen, intestine, gill, muscle and skin the uptake showed the opposite behavior. The small percentage of uptake in the brain can be attributed to the blood-brain barrier (BBB) that prevents molecules from reaching the brain (Shakeri *et al.*, 2020). Free  $^{67}\text{GaCl}_3$  showed a much more general biodistribution to most organs, while also showing elevated uptake in

the liver ( $8.51 \pm 1.45$  % ID/g at 24 h,  $11.8 \pm 3.02$  % ID/g at 48 h after injections), kidney ( $6.01 \pm 2.47$  % ID/g at 24 h,  $5.38 \pm 1.15$  % ID/g at 48 h post injection), and spleen ( $6.53 \pm 1.99$  % ID/g

at 24 h after injections) in rainbow trout (Fig. 4).




Figure 1: TEM image of gamma-irradiated (at a dose of 30 KGy) RE-NPs were immobilized on a coated copper grid at room temperature (n=9).






Figure 2: RTLC of (A)  $^{67}\text{Ga}$ -RE-NPs (1%, 30 KGy) and (B) Free  $^{67}\text{GaCl}_3$  in Sodium acetate and acetic acid as mobile phase (n=9).



**Figure 3:** Organ concentrations of radioactivity (presented as SUV, mean  $\pm$  SD) obtained at 24 h and 48 h after the injections of  $^{67}\text{Ga}$ - RE-NPs and free  $^{67}\text{GaCl}_3$  (n = 9), error bars represent standard deviation.



**Figure 4:** Representative SPECT whole-body images of  $^{67}\text{Ga-RE-NPs}$  and free  $^{67}\text{GaCl}_3$  biodistribution in healthy rainbow trout 24 h (A) and 48 h (B) after intraperitoneal injection (n=9). Organs with significant uptake are labeled.

## Discussion

This study demonstrated the feasibility and validity of SPECT imaging to investigate tissue accumulation of RE-NPs in healthy rainbow trout. SPECT is a powerful and non-invasive imaging technique to visualize the biodistribution of molecules labeled with radioactive isotopes such as  $^{67}\text{Ga}$ ,  $^{99\text{m}}\text{Tc}$  and  $^{123}\text{I}$  (Knight *et al.*, 2019). Different parameters affect the labeling reaction; such as the pH of the reaction medium, Rosmarinic acid concentration, and reaction time and temperature (El-Sharawy *et al.*, 2022). This current study shows the radiolabeled RE-NPs at 1% concentration were prepared using  $^{67}\text{Ga}$  with good labeling at about 97% efficiency (RTLC method). El-Sharawy *et al.* (2022) also reported radiolabeling yield of Rosmarinic acid with  $^{99\text{m}}\text{Tc}$  radioisotope was 92.2% in pH=4 for 15 min.

To the best of our knowledge, this is the first report of SPECT imaging in rainbow trout demonstrating the feasibility of tracking of RE-NPs uptake. Heidarieh *et al.* (2014) injected  $^{67}\text{Ga}$ -alginate nanoparticles into rainbow trout through a peritoneum to determine the whole-body distribution of alginate nanoparticles. In this study, the biodistribution of  $^{67}\text{Ga}$ -RE-NPs was performed to assess their behavior as potential natural antioxidant agents. The imaging technique using SPECT camera detected higher net uptake in the kidney after 24 h post injection of  $^{67}\text{Ga}$ -RE-NPs in healthy rainbow trout.

Cells have crucial antioxidant defense mechanisms to protect themselves

against toxic injury by free-radicals. The enzymatic antioxidants are SOD, CAT, and GPx (Pisoschi and Pop, 2015; Mirończuk-Chodakowska *et al.*, 2018). SOD<sub>1</sub> accounts for the highest proportion of SOD activity in the kidney (Schieber and Chandel, 2014). SOD<sub>2</sub> is also expressed in most tissue cells, such as the stomach, skeletal muscle, spleen, heart, liver, and the brain (Van Remmen *et al.*, 1999). SOD<sub>3</sub> is highly expressed in blood vessels, kidneys and the heart (Wang *et al.*, 2018). CAT is abundantly present in the liver and kidneys (Ho *et al.*, 2004). The kidney is a major source of plasma GPx (Avissar *et al.*, 1994). Based on recent published data, plant extracts enriched with polyphenols could improve the enzymatic antioxidant defense system in salmon (Santana *et al.*, 2021). Among natural antioxidant plants, rosemary was known for the presence of phenolic compounds (Collins and Charles, 1987; Moreno *et al.*, 2006). Based on recently research findings, biodistribution data and increased renal and splenic uptake of  $^{67}\text{Ga}$ -RE-NPs in fish after IP administration can indicate that RE-NPs can improve the renal and splenic enzymatic antioxidant defense system in rainbow trout. The present results also were confirmed by the work presented by Hernández *et al.* (2014) and Rezanejad *et al.* (2019b) indicating that using natural antioxidant RE-NPs in the diet can improve the resistance of fish muscle against oxidation and also fish fillet quality. Thus, based on the above investigation it is evident firstly that the potential of  $^{67}\text{Ga}$ -RE-NPs to determine

RE-NPs pharmacokinetics, elimination pathway, metabolism, and potential degradation, shows *in vivo* safety in rainbow trout. Secondly, the supplemented feed with gamma irradiated RE-NPs can be a good candidate for modulating the rainbow trout antioxidant system activities.

The applications of nanoparticles are widely studied for both treatment applications and molecular imaging. In this study, the radiolabeled RE-NPs at 1% concentration were prepared using <sup>67</sup>Ga with good labeling at about 97% efficiency (RTLC method). Biodistribution of the radiolabeled RE-NPs was checked in healthy rainbow trout up to 24 h compared to free <sup>67</sup>GaCl<sub>3</sub>. The data showed that the tracer accumulation was high in kidney and spleen tissues. Based on the current study, the radiolabeling of RE-NPs as natural tracers can be a true and sensitive method to image and understand their whole-body distribution and pharmacokinetics in fish models.

## References

- Amelinckx, S. and Van Landuyt, J., 2003.** Transmission Electron Microscopy. In: Encyclopedia of Physical Science and Technology, 3rd Edition, Robert Meyers A., editor, Academic Press, 53–87, Cambridge, Massachusetts, USA.
- Avissar, N., Ornt, D.B., Yagil, Y., Horowitz, S., Watkins, R.H., Kerl, E.A., Takahashi, K., Palmer, I.S. and Cohen, H.J., 1994.** Human kidney proximal tubules are the main source of plasma glutathione peroxidase. *American Journal of Physiology - Cell Physiology*, 266, 367–375. <https://doi.org/10.1152/ajpccell.1994.266.2.c367>.
- Boyer, C., Whittaker, M.R., Bulmus, V., Liu, J.Q. and Davis, T.P., 2010.** The design and utility of polymer-stabilized iron-oxide nanoparticles for nanomedicine applications. *NPG Asia Materials*, 2(1), 23–30. <https://doi.org/10.1038/asiamat.2010.6>.
- Collins, M.A. and Charles, H.P., 1987.** Antimicrobial activity of Carnosol and Ursolic acid: two anti-oxidant constituents of *Rosmarinus officinalis* L. *Food Microbiology*, 4(4), 311–315. [https://doi.org/10.1016/s0704-0020\(87\)80005-9](https://doi.org/10.1016/s0704-0020(87)80005-9).
- El-Sharawy, D.M., Khater, S.I., El Refaye, M.S., Hassan, H.M., AboulMagd, A.M. and Aboseada, M.A., 2022.** Radiolabeling, biological distribution, docking and ADME studies of <sup>99m</sup>Tc-Ros as a promising natural tumor tracer. *Applied Radiation and Isotopes: Including Data, Instrumentation and Methods for use in Agriculture, Industry and Medicine*, 184, 110196.
- Heidarieh, M., Daryalal, F., Mirvaghefi, A., Rajabifar, S., Diallo, A., Sadeghi, M., Zeiai, F., Moodi, S., Maadi, E., Sheikhzadeh, N., Heidarieh, H. and Hedyati, M., 2014.** Preparation and anatomical distribution study of <sup>67</sup>Ga-alginate nanoparticles for SPECT purposes in rainbow trout (*Oncorhynchus mykiss*). *Nukleonika*, 59(4), 153–159.

- <https://doi.org/10.2478/nuka-2014-0019>.
- Hernández, A., García, B.G., Jordán, M.J. and Hernández, M.D., 2014.** Improved conservation of gilthead seabream (*Sparus aurata*) in ice storage. The influence of doses of rosemary extract added to feed. *Aquaculture*, 426–427, 31–40. <https://doi.org/10.1016/j.aquaculture.2014.01.018>.
- Ho, Y.S., Xiong, Y., Ma, W., Spector, A. and Ho, D.S., 2004.** Mice lacking catalase develop normally but show differential sensitivity to oxidant tissue injury. *Journal of Biological Chemistry*, 279(31), 32804–32812. <https://doi.org/10.1074/jbc.m404800200>.
- Jalilian, A.R., Mirsadegh, L., Haji-Hosseini, R., Rajabifar, S. and Bolurinovin, F., 2007.** Production and evaluation of [<sup>67</sup>Ga]-DTPA-Rituximab. *Iranian Journal of Radiation Research*, 4(4), 187–193.
- Jalilian, A.R., Hosseini, M.A. and Bolourinovin, F., 2011.** Preparation and biodistribution study of <sup>67</sup>Ga-gallium calcitonin. *Iranian Journal of Nuclear Medicine*, 19(2), 38–45.
- Kitson, S.L., Moody, T. and Watters, W., 2014.** *Modern Developments in Isotopic Labelling*. Almac Group, Northern Ireland, UK, 11p.
- Knight, J.C., Mosley, M.J., Dias, Kersemans, V., Dias, G.M., Allen, P.D., Smart S. and Cornelissen, B., 2019.** Dual-isotope imaging allows in vivo immunohistochemistry using radiolabelled antibodies in tumours. *Nuclear Medicine and Biology*, 70, 14–22.
- <https://doi.org/10.1016/j.nucmedbio.2019.01.010>.
- Madsen, H.L. and Bertelsen, G., 1995.** Spices as antioxidants. *Trends in Food Science and Technology*, 6(8), 271–278. [https://doi.org/10.1016/s0924-2244\(00\)89112-8](https://doi.org/10.1016/s0924-2244(00)89112-8).
- Mirończuk-Chodakowska, I., Witkowska, A.M. and Zujko, M.E., 2018.** Endogenous non-enzymatic antioxidants in the human body. *Advances in Medical Sciences*, 63(1), 68–78. <https://doi.org/10.1016/j.advms.2017.05.005>.
- Moreno, S., Scheyer, T., Romano, C.S. and Vojnov, A.A., 2006.** Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. *Free Radical Research*, 40(2), 223–231. <https://doi.org/10.1080/10715760500473834>.
- Offord, E.A., Guillot, F., Aeschbach, R., Lölliger, J., and Pfeifer, A.M.A. 1997.** Antioxidant and biological properties of rosemary components: implications for food and health. In: *Natural antioxidants: chemistry, health effects, and applications*, Shahidi F. editor, AOCS Press, Champaign, Illinois, USA, 88–96.
- Orlando, P., Binaglia, L., De Feo, A., Trevisi, R., Melodia, C. and Trenta, R., 1994.** Preparation of high molecular weight radioiodinated alginic acid. *Journal of Labelled Compounds and Radiopharmaceuticals*, 34(7), 653–

657.  
<https://doi.org/10.1002/jlcr.2580340>  
709.
- Pintore, G., Usai, M., Bradesi, P., Juliano, C., Boatto, G., Tomi, F., Chessa, M., Cerri, R., and Casanova, J., 2002.** Chemical composition and antimicrobial activity of *Rosmarinus officinalis* L. oils from Sardinia and Corsica. *Flavour and Fragrance Journal*, 17(1), 15–19.  
<https://doi.org/10.1002/ffj.1022>.
- Pisoschi, A.M. and Pop, A., 2015.** The role of antioxidants in the chemistry of oxidative stress: a review. *European Journal of Medicinal Chemistry*, 97, 55–74.  
<https://doi.org/10.1016/j.ejmech.2015.04.040>.
- Rezanejad, R., Ojagh, S.M., Heidarieh, M., Raeisi, M., Rafiee, G. and Alishahi, A., 2019a.** Characterization of gamma-irradiated Rosemary (*Rosmarinus officinalis*). *Turkish Journal of Pharmaceutical Sciences*, 16(1), 43–47.  
<https://doi.org/10.4274/tjps.37880>.
- Rezanejad, R., Ojagh, S.M., Heidarieh, M., Raeisi, M., Alishahi, A. and Rafiee, G., 2019b.** The impact of diets supplemented with different forms of Rosemary and BHA on chemical, microbial and sensory properties of rainbow trout fillet. *Journal of Aquatic Food Product Technology*, 28(5), 478–494.  
<https://doi.org/10.1080/10498850.2019.1604594>.
- Saha, G.B., 2004.** *Fundamentals of Nuclear Pharmacy*, 5th edition. Springer-Verlag, New York, USA.
- Santana, P.A., Jara-Gutiérrez, C., Mellado, M., Carlos Forero, J., Guzmán, F., Barriga, A., Albericio, F., and Álvarez, C.A., 2021.** Effects of elderflower extract enriched with polyphenols on antioxidant defense of salmon leukocytes. *Electronic Journal of Biotechnology*, 52, 13–20.  
<https://doi.org/10.1016/j.ejbt.2021.04.004>.
- Schieber, M. and Chandel, N.S., 2014.** ROS function in redox signaling and oxidative stress. *Current Biology*, 24(10), 453–462.  
<https://doi.org/10.1016/j.cub.2014.03.034>.
- Shahidi, F., 1997.** Natural antioxidants: an overview. In: *Natural antioxidants: chemistry, health effects, and applications*, Shahidi F. editor, AOCS Press, Champaign, Illinois, USA, 1–11.
- Shakeri, S., Ashrafizadeh, M., Zarrabi, A., Roghanian, R., Ghasemipour Afshar, E., Pardakhty, A., Mohammadinejad R., Kumar A. and Kumar Thakur, V., 2020.** Multifunctional polymeric nanoplateforms for brain diseases diagnosis, therapy and theranostics. *Biomedicines*, 8(1), 13.  
<https://doi.org/10.3390/biomedicines8010013>.
- Tironi, V.A., Tomás, M.C. and Añón, M.C., 2010.** Quality loss during the frozen storage of sea salmon. Effect of rosemary (*Rosmarinus officinalis*) extract. *LWT – Food Science and*

- Technology*, 43(2), 263–272.  
<https://doi.org/10.1016/j.lwt.2009.07.007>.
- Van Remmen, H., Salvador, C., Yang, H., Huang, T.T., Epstein, C.J. and Richardson, A., 1999.** Characterization of the antioxidant status of the Heterozygous Manganese Superoxide Dismutase Knockout mouse. *Archives of Biochemistry and Biophysics*, 363(1), 91–97.  
<https://doi.org/10.1006/abbi.1998.1060>.
- Wang, Y., Branicky, R., Noë, A. and Hekimi, S., 2018.** Superoxide dismutases: dual roles in controlling ROS damage and regulating ROS signaling. *Journal of Cell Biology*, 217(6), 1915–1928.  
<https://doi.org/10.1083/jcb.201708007>.
- Xie, J., Van Alstyne, P., Uhlir, A. and Yang, X., 2017.** A review on rosemary as a natural antioxidation solution. *European Journal of Lipid Science and Technology*, 119(6), 1600439.  
<https://doi.org/10.1002/eilt.201600439>.